

# PSYCHE

VOL. XXI.

DECEMBER, 1914.

No. 6

## SOME MYRMECOPHILOUS INSECTS FROM MEXICO.<sup>1</sup>

BY W. M. MANN.

The following notes are based on a small collection made during the months of May, June and July, 1913, in the State of Hidalgo, Mexico. Most of this time was spent at the Guerrero Mill, located below Real del Monte, and at the Hacienda de Velasco, where I was the guest of Mr. and Mrs. C. W. Van Law. They, together with the other Americans and English in charge of the mines and mills, showed me every hospitality and it is largely due to them that I was able to reside and collect in this interesting locality. I am much indebted to Miss Helen L. Locke, Mr. and Mrs. J. H. Skews and Messrs. Broiderick, Benton, Funston and Calland, not only for assistance of various kinds, but for many desirable specimens which they collected. The material collected has been placed in the collections of the Museum of Comparative Zoölogy at Cambridge and of Mr. B. Preston Clark of Boston, through the kindness of whom the excursion was possible.

Although this region is easily accessible, and unusually interesting and rich in insect life because of its ecologically varied nature, it has been largely neglected by collectors. Only a few things have been collected at Pachuca and at Guerrero Mill. Prof. W. M. Wheeler, in looking over the literature on Mexican ants, failed to find Hidalgo cited as the locality for a single species, and I have found no records of other insects from that state.

Among the thirty-nine species and varieties of ants collected (an account of these has been published by Wheeler in the *Journal of the New York Entomological Society*, Vol. XXII, 1914, pp. 37-61) nine were found to harbor guests or parasites. Three of these belong to the genus *Formica*, one each to *Camponotus*, *Prenolepis*,

<sup>1</sup> Contribution from the Entomological Laboratory of the Bussey Institution of Harvard University, No. 84.

Liometopum and Tapinoma, and two to Pheidole, all of which genera are commonly the hosts of inquilines, Prenolepis less frequently than the others. The inquilines found are mostly undescribed species of genera known to be of myrmecophilous habits, but a new Diaprid, *Hemilexis jessei*, and a new genus of Bethyridæ, *Bruesiella formicaria*, both of which are probably parasitic on the host ant, are described. There is also an interesting new genus of cockroach, *Myrmecoblatta rehni*, which inhabits nests of various species of ants, where it no doubt officiates as scavenger. The occurrence of the weevil *Liometophilus manni* Fall at Pachuca, with the same species of host with which it has been found in Arizona shows that it is definitely associated with this ant.

I have included in this paper a few remarks on the synonymy of some species from north of the Mexican boundary, and erected a new subgenus to contain certain of our species of *Cremastocheilus*.

## ORTHOPTERA.

Family BLATTIDÆ.

Subfamily BLATTINÆ.

*Myrmecoblatta* gen. nov.

*Female*: Body elongate oval, not greatly flattened, the abdomen about as broad as the thorax. Head almost concealed by the pronotum, only a narrow strip of the vertex showing. Head small, triangular, the vertex somewhat swollen. Eyes well developed, elongate, emarginate on inner border. Antennæ long and slender, the first joint thickened, the second small, the third longer than the second and twice the length of the fourth. Thorax longer than broad, convex above, tegmina and wings absent. Abdomen convex above, beneath gradually elevated to the middle, which is therefore broadly carinate. Supra-anal plate semi-orbicular, evenly rounded in outline. Cerci large, with eight distinct joints. Sub-genital plate apparently provided with valves. Last ventral segment completely divided by a V-shaped slit. Legs short and robust, with abundant fine, short hairs. Femora weakly armed beneath, and with a strong spine at apex; tibiæ spined at apex and on outer margin.

*Male*: Form broad. Pronotum entirely concealing the head. Tegmina more than half as broad as long; short, not reaching apex of abdomen, with poorly developed veins. Wings short and broad, less than half the length of tegmina, with poorly developed veins. Supra-anal plate more than twice as long as broad, the sides concave in outline, posterior border notched at middle, rounded on either side. Sub-genital plate broader than long, straight at sides, rounded behind, slightly concave at middle, bearing two equal stylets.

*Type*: *Myrmecoblatta rehni* sp. nov.

***Myrmecoblatta rehni* sp. nov.**

*Female:* (Fig. 1). Length 5 mm. Width 3 mm. Color reddish-brown, the legs lighter, above regularly punctate and finely setose. Thorax longer than broad. Pronotum broader than long, with concave posterior border and extended, narrowly rounded angles.

Meso- and epinota subequal, with nearly straight sides and concave posterior border. Abdominal segments 1-7 subequal, with straight posterior borders, the eighth considerably narrower than the seventh, rounded behind. Cerci 8-jointed, a third as broad as long, acuminate for apical half their length; nearly as long as the sub-genital plate. Head about as broad as long, the front with an indistinct Y-shaped suture. Labium obtusely pointed, concealing the mandibles. Eyes occupying the sides of the head, faintly emarginate on the inner border. Antennæ slender, shorter than the body, about 37-jointed; the third joint twice as long as the fourth, joints 4-13 transverse, the rest distinctly longer than broad, and sub equal; all the joints finely setose. Legs short, rather flat,

the femora armed at apex with a slender curved spine; tibiæ bearing on the outer border three slender subequal spines, at the apex two long and two shorter spines.

*Male:* (Fig. 2). Length 4.5 mm. Width 2 mm. Color and setosity similar to

that of female. Pronotum proportionally slightly broader than in the female, the angles broadly rounded, posterior border concave. Tegmina extending a little more than two-thirds the length of abdomen, broadly oval in outline, the apex narrowly rounded; venation not very distinct. Wings less than half the length of tegmina. Cerci




Fig. 1. *Myrmecoblatta rehni* gen. & sp. nov.  
Ventral surface of female.




Fig. 2. *Myrmecoblatta rehni* gen. & sp. nov.  
Ventral surface of last gastric segments of male.

more slender than in the female, longer than the supra-anal plate at middle. Legs and antennæ longer than in the female, the latter as long as the body.

*Nymph*: Similar to female, lighter in color. The suture on the front of head is more distinct than in the adult.

Described from one male and numerous females and nymphs taken at Guerrero Mill in nests of *Formica subcyanea* Wheeler, *F. rufibarbis* Fab. var. *gnava* Buckley, and *Camponotus maculatus* Fab. subsp. *picipes* Olivier. Some of the females had oöthecæ attached. They were abundant, several occurring in almost every nest, where they are no doubt very efficient scavengers. The genus has no systematic affinities with *Attaphila*, the only other known Blattid of Myrmecophilous habits, but is more closely related to *Blatta*, to which it runs in Caudell's key to the genera of this family (*Proc. U. S. Nat. Mus.* Vol. 44, 1913, p. 601-602).

## COLEOPTERA.

### Family STAPHYLINIDÆ.

#### *Pseudolomechusa* subgen. nov.

Type: *Xenodusa sharpi* Wasmann.

This beetle was described by Wasmann from a series which were taken at Cuernavaca, in company with *Camponotus auricomus* Roger. In one canyon near the Guerrero Mill I found a number of adults, some of them in copula, with colonies of *Camponotus maculatus* var. *picipes*, Olivier which was here very abundant. In the immediate vicinity were many nests of *Formica microgyna rasilis* var. *nahua* Wheeler and several of *T. rufibarbis* var. *gnava*. Buckley. With the former ant three adults of *sharpi* were taken and with *gnava* several more, and three larvæ, the latter close to the ant brood. As most of the beetles were found with *C. picipes* (nineteen were taken from one nest) it appears that this is the secondary or winter host and *F. gnava* the primary or definitive. This heterocious habit is common to all of our species of *Xenodusa* as far as known. The adult *sharpi* is very different from the other American species on account of its greatly thickened pronotal margins and shorter antennal joints. Seen in profile the anterior border is drawn out beneath into a distinct angle similar to the structure in the species of the European genus *Lomechusa*.

The larva is very different from that of *X. cava*, the only one known in that stage, which has been described and figured by Wheeler (*Jour. N. Y. Ent. Soc.*, Vol. XIX, 1911, p. 165-166) and closely resembles that of *Lomechusa*. Those which I found are 5.5 mm. in length, yellowish-white in color, with short, feeble legs, no eyes, papillose antennæ and shallowly impressed vertex. The body is slender and cylindrical, not at all depressed as it is in *cava*, with an even covering of fine erect hairs.

The movements in life were very slow. Considering the pronotal structure and the larval characters *sharpi* should not be included in *Xenodusa* sens. strict., but more properly belongs to a distinct subgenus for which I propose the name *Pseudolomechusa*.

*Apteronina wasmanni* sp. nov.

Length 3 mm. Head broader than long, rounded at sides, occipital margin straight; clypeus projecting, the anterior border rounded; front broadly depressed, the surface concave, with a longitudinal impression extending to occiput. Eyes rather large, located at sides of the anterior third of head. Antennæ long, the first joint about twice the length of the second, which is two-thirds as long as the third; the second joint is sharply constricted near base; fourth joint about half as long as third; joints 4-10 subequal in length, becoming thicker anteriorly; apical joint cylindrical, nearly twice the length of penultimate. Thorax not as broad as head, very slightly longer than broad, rounded in front and at anterior third of sides, posterior to which the sides are straight; posterior border straight; surface slightly concave, with a narrow longitudinal impression at middle. Elytra transverse, each as long as broad, together broader than thorax; sides nearly straight; posterior border rounded. Abdomen at base as broad as elytra, broadest behind. Legs long and slender.

Body and legs subopaque, finely punctate, the punctures coarser on the elytra and abdomen; everywhere with fine, recumbent, silky hairs; thorax and sides of abdomen with longer coarse hairs.

Color black, the antennæ and legs brownish.

Described from several specimens taken with *Liometopum apiculatum* Mayr at Pachuca and Guerrero Mill.

*A. schmitti* Wasmann, which lives in company with the same host ant from Colorado to Southern Arizona, has the head longer than broad, the thorax is longer in proportion to the width, the general form is more slender; the color is light ferruginous, except the gaster, which is dark fuscous. The color of the Mexican form of the host ant is considerably darker than that of the Arizona variety with which I have taken *schmitti*, so the beetle in each



case resembles the ant in that respect. Besides color the beetle is similar to the host in pilosity and, superficially, in form, and in life it carries its abdomen erect in a manner similar to the way in which the ant holds its gaster. Nothing is known regarding the biological relation of the beetle to the ant.

*Dinardella mexicana* sp. nov.

Length 2.75 mm. Head slightly broader than long, rounded at sides and in front, posterior border straight; front with a flattened disc. Eyes medium in size, flat. Antennæ short, extending to apex of elytra; joints in front of the middle thickest; first joint as long as the two succeeding together; joints 2-3 longer than broad; joints 4-10 transverse; apical joint a little less than twice the length of penultimate. Prothorax as long as broad, broadest at posterior fourth, posterior to which the sides are straight; anterior and posterior borders straight; disc with a faint longitudinal impression. Elytra transverse at base, together considerably broader than thorax; sides slightly rounded; the posterior border of each straight. Abdomen broad and flat; at base as broad as elytra, the greatest width at the third and fourth segments. Legs short and slender.

Body subshining, finely punctate throughout and covered with a fine mat of yellow pubescence; the thorax and elytra bear scattered, erect, black hairs.

Color dark fuscous to piceous, legs and antennæ lighter.

Described from a small series taken at Pachuca and Guerrero Mill with *Liometopum apiculatum* Mayr.

Besides being much darker in color *D. mexicana* differs from *D. liometopi* Wasmann in the shape of the head, which in the latter species is distinctly triangular and much broader than long; in the thorax not being broader than long, and in having proportionately longer elytra.

*Zyras (Myrmecia) tapinomatis* sp. nov.

Length 3 mm. Thorax broad; abdomen narrow. Color piceous, except the antennæ, mandibles, palpi and tarsi, which are fuscous; shining throughout; the antennæ less so than the body. Head not as wide as prothorax, about as long as broad; finely but distinctly punctate. Antennæ not extending to apex of elytra, thick; first and third joints longer than broad, second joint small; joints 4-10 as long as broad, each slightly longer than the preceding; apical joint a little shorter than the next two together. Pronotum a little broader than long, with rounded sides, the disc at middle very broadly and deeply impressed, the impression about one and a half times as long as broad; finely punctate. Elytra together broader than long, the sides straight and parallel, punctured similar to pronotum. Abdomen long and slender; about two-thirds as broad as elytra; very shining; the third and fourth segments each having on the dorsum a prominent tubercle, the anterior of which is the smallest, conical, and bears at the sides a fine mat of re-

cumbent yellow hairs; the posterior from above is round, with flat surface and without hairs. The median portion of the abdominal dorsum is glabrous; the rest of the body, legs and antennæ are thickly though microscopically pilose, the antennæ more densely than the other parts.

Described from three specimens taken at San Miguel in nests of *Tapinoma sessile* Say\*.

This species is very similar to *M. lugubris* Casey which I took some years ago in Orcus Island (Puget Sound) with colonies of the same host ant, but differs in the fine punctation of the head and thorax, which in *lugubris* are reticulate. The thoracic depression in *tapinomatis* is deeper and narrower than in *lugubris* and the abdomen is proportionally much more slender.

#### Family PSELAPHIDÆ.

##### *Pilopius major* sp. nov.

*Female*: Length 3.25 mm. Color throughout brown. Head narrower than thorax; from the front twice as long as broad, the sides in front of eyes sub-parallel; vertex foveolately impressed at sides, a thin median carina extending to between the antennal tubercles. Eyes large and convex, located at the posterior third of head. Antennæ thick, the first joint slightly bent at base, as long as the two succeeding joints together; third joint smaller than the fourth, which is larger than the fifth; joints 5-11 sub-globose, equal in size; apical joint one and three-fourths times as long as penultimate, thickened. Prothorax a little longer than broad, narrowed in front, with rounded sides; base compressed above, with median and lateral foveæ; the former is elongate, extending one-third the distance to apex. Elytra together longer than broad, the sides rounded; humeri elevated into thick ridges; discal striæ broad, slightly arcuate, extending three-fourths the length of elytra. Abdomen narrower than elytra and about the same length.

Body shining everywhere, finely punctate, with a thin covering of short, scale-like hairs which are more dense in the pronatal foveæ and on the occiput and posterior elytral margins.

Described from specimens taken at San Miguel with *Prenolepis* (*Nylanderia*) *mexicana* Forel., one from each of three colonies. The maxillary palpi are short and proportionally smaller than in the other species of the genus that I have seen. This species belongs to the group which includes *pulvereus* Lec., *ocularis* Csy. and *abruptus* Csy., in the latter two of which only the males have been described, but is considerably larger than any of these. The pubescence is much sparser than in *pulverius*. The other species whose hosts are known are guests of different species of Aphæno-

gaster. Like most other Pselaphids of myrmecophilous habits, few are found in a single ant colony, but they are generally common, there being one or two in almost every nest of the host ant in the localities where they occur.

### Family HISTERIDÆ.

#### *Hetærius helenæ* sp. nov.

Length 2 mm. Color reddish-brown, in some specimens the side margins of the head, the margin of mandible and apex of femur piceous; shining. Form robust, about two-thirds as broad as long. Head above sparsely, coarsely punctate, less so between the eyes and beneath; clypeus about twice as broad as long, finely punctate; front with short yellow hairs. Thorax transverse, its breadth nearly twice the length; broadest at base; front border concave, narrowly margined; sides straight to the constriction, which is located about five-eighths the distance from the anterior angle to base; median portion of disc finely, regularly punctate, with sparse yellow hairs; marginal portion in front of constriction very finely rugose longitudinally and less shining than the remainder of thorax; the anterior border feebly rounded, posterior portion slightly elevated; margin of sides with a rather dense brush of long stiff yellow hairs. Elytra at base distinctly broader than thorax, the width of both together about equal to length; abundantly pilose; apical margin punctate; marginal and humeral striæ entire; third and fourth striæ extending for five-sixths the length of elytra, carinated for entire length. Propygidium and pygidium finely rugulose, the former with sparse long hairs. Posternum narrowly margined for half its length; transversely impressed at half the distance from end of margination to anterior border; distinctly excised at anterior border; margined portion shining, smooth behind and finely punctate in front. Anterior tibiæ nearly a third as broad as long, the front margin posterior to the angle straight, with about eleven elongate denticles. Middle and posterior tibiæ one-third as broad as long, the outer edge rounded.

There is considerable variation in the shade of color and a little in size. The hairs are rather long, curved, and moderately abundant. In some specimens they are almost entirely absent, probably having been scraped off by the ants.

Described from a number of specimens taken in nests of *Formica subcyanea* Wheeler and *F. microgyna* subsp. *rasilis* var. *nahua* Wheeler. The former ant is the favorite host, as nearly every nest in some localities contained the beetles, while they were found in only one nest of the latter species out of dozens examined.

This species, the first known from Mexico, is more closely related to some undescribed forms from our southwestern states than to any of the described species. It is more pilose than any of the species which I have seen, except *californicus* and an undescribed species (*hirsutus* mss.) from Arizona.



*Terapus mnizechi* Mars.

Two specimens were taken at San Miguel in nests of *Pheidole vasleti* var. *acolhua* Wheeler. In the place where these were found was a stone fence extending for about a half mile, along which were literally hundreds of nests of *acolhua*. Although I made several visits here and searched many nests in other localities for the beetles no more were found. It is evidently exceedingly rare locally but has a wide distribution in Mexico.

*Terapus infernalis* Fall.*Melanetarius infernalis* Fall.

An account of this species has been given by Wheeler (PSYCHE XVIII, 1911, p. 112-114) who found a number of specimens at Pasadena in company with *Pheidole hyatti* Emery. One of these was studied by Bickhardt (PSYCHE XIX, p. 97, 1912) who found that the beetle and most of the details of Wheeler's figure agreed with Marseul's description and figure of *mnizechi*, to which species he referred it. Recently, through the kindness of Mr. J. H. Arrow, I was able to examine in the British Museum a specimen of *mnizechi* from the Federal District of Mexico. This was undoubtedly the same as the San Miguel specimens, and these are specifically quite distinct from the Southern California form. In *infernalis* the propygidium is feebly punctate above and more coarsely beneath, with the upper, smoother portion much smaller than the lower, and there is no sharp line between the two parts. In *mnizechi* the upper portion is smooth and shining, the lower punctured and opaque, and each part is in strong contrast to the

Correction: For *Myrmecotonus* read "*Myrmeceicon*," page 179 and 180.

mentum angulate behind. This character is correlated with the geographical distribution of the species, all of the species in this group inhabiting the middle and far West. *C. wheeleri* Lec.

gaster. Like most other Pselaphids of myrmecophilous habits, few are found in a single ant colony, but they are generally common, there being one or two in almost every nest of the host ant in the localities where they occur.

### Family HISTERIDÆ.

#### *Hetærius helenæ* sp. nov.

Length 2 mm. Color reddish-brown, in some specimens the side margins of the head, the margin of mandible and apex of femur piceous; shining. Form robust, about two-thirds as broad as long. Head above sparsely, coarsely punctate, less so between the eyes and beneath; clypeus about twice as broad as long, finely punctate; front with short yellow hairs. Thorax transverse, its breadth nearly twice the length; broadest at base; front border concave, narrowly margined; sides straight to the constriction, which is located about five-eighths the distance from the anterior angle to base; median portion of disc finely, regularly punctate, with sparse yellow hairs; marginal portion in front of constriction very finely rugose longitudinally and less shining than the remainder of thorax; the anterior border feebly rounded, posterior portion slightly elevated; margin of sides with a rather dense brush of long stiff yellow hairs. Elytra at base distinctly broader than thorax, the width of both together about equal to length; abundantly pilose; apical margin punctate; marginal and humeral striæ entire; third and fourth striæ extending for five-sixths the length of elytra, carinated for entire length. Propygidium and pygidium finely rugulose, the former with sparse long hairs. Posternum narrowly margined for half its length; transversely impressed at half the distance from end of margination to anterior border; distinctly excised at anterior border; margined portion shining, smooth behind and finely punctate in front. Anterior tibiæ nearly a third as broad as long, the front margin posterior to the angle straight, with about eleven elongate denticles. Middle and posterior tibiæ one-third as broad as long, the outer edge rounded.

There is considerable variation in the shade of color and a little in size. The hairs are rather long, curved, and moderately abundant. In some specimens they are almost entirely absent, probably having been scraped off by the ant.

More than any other I have seen, except *californicus* and an undescribed species (*hirsutus* mss.) from Arizona.

*Terapus mnizechi* Mars.

Two specimens were taken at San Miguel in nests of *Pheidole vasleti* var. *acolhua* Wheeler. In the place where these were found was a stone fence extending for about a half mile, along which were literally hundreds of nests of *acolhua*. Although I made several visits here and searched many nests in other localities for the beetles no more were found. It is evidently exceedingly rare locally but has a wide distribution in Mexico.

*Terapus infernalis* Fall.*Melanetarius infernalis* Fall.

An account of this species has been given by Wheeler (PSYCHE XVIII, 1911, p. 112-114) who found a number of specimens at Pasadena in company with *Pheidole hyatti* Emery. One of these was studied by Bickhardt (PSYCHE XIX, p. 97, 1912) who found that the beetle and most of the details of Wheeler's figure agreed with Marseul's description and figure of *mnizechi*, to which species he referred it. Recently, through the kindness of Mr. J. H. Arrow, I was able to examine in the British Museum a specimen of *mnizechi* from the Federal District of Mexico. This was undoubtedly the same as the San Miguel specimens, and these are specifically quite distinct from the Southern California form. In *infernalis* the propygidium is feebly punctate above and more coarsely beneath, with the upper, smoother portion much smaller than the lower, and there is no sharp line between the two parts. In *mnizechi* the upper portion is smooth and shining, the lower punctured and opaque, and each part is in strong contrast to the other. The upper part is proportionally larger than it is in *infernalis*. The latter species is smaller, measuring two millimeters while *mnizechi* measures three.

## Family SCARABÆIDÆ.

*Creastocheilus*, subgenus *Myrmecotonus* nov.

Type: *Creastocheilus knochii* Lec.

I propose this subgenus to contain those species which have the mentum angulate behind. This character is correlated with the geographical distribution of the species, all of the species in this group inhabiting the middle and far West. *C. wheeleri* Lec.

should be included with these, though in some specimens the mentum is very feebly notched.

The species which belong to *Cremastocheilus* subgenus *Cremastocheilus* have the mentum distinctly incised behind and are all native of the Atlantic states. The type of the genus *Cremastocheilus* is *C. castanea* Knoch.

*Cremastocheilus* (*Myrmecotonus*) *mexicanus* West.

Several specimens, found with *Formica subcyanea* Wheeler at Guerrero Mill agree closely with others identified as this species from nests of *Formica gnava* Buckley in the Huachuca Mts. (Schaeffer) and from the Santa Rita Mts., Arizona (Snow). There is in this series, as in other species in the genus, considerable variation in sculpture and pilosity as well as in size.

*Cremastocheilus* (*Myrmecotonus*) *armatus* Walker.

*C. pilisicollis* Horn.

Recently I examined the type and several paratypes of *C. armatus* in the British Museum, and compared with specimens of the form which is identified in collections as *pilisicollis* Horn. The two are identical, so the latter name must fall into synonymy, as was considered probable by Horn. (Monographic Revision of the species of *Cremastocheilus* and Synopsis of the Euphoriae of the United States. *Proc. Amer. Phil. Soc.* XVIII, 1879, p. 390).

Family CURCULIONIDÆ.

*Liometophilus manni* Fall.

A single specimen of this curious weevil was taken at Pachuca in the galleries of a colony of *Liometopum apiculatum* Mayr. The species was first found in Southern Arizona, where it lives with a variety of the same ant. The oily-red color, so characteristic of the chitin in many beetles of myrmecophilous habits, is evident at the middle of the anterior elytral margin, where the scales are absent, probably having been gnawed off by the ants. This is, so far as is known, the only truly myrmecophilous weevil found in North America.

The Mexican specimen (length 4 mm.) is considerably larger than a paratype from Arizona in my collection, and the light-

colored transverse bands on the elytra are in stronger contrast to the rest of the body, but otherwise the two agree closely.

## HYMENOPTERA.

### Family DIAPRIDÆ.

#### *Hemilexis jessei* sp. nov.

*Female:* (Fig. 3). Length 3 mm. Color jet black, except the mandibles, front of the propleuræ, petiole, antennæ and legs, which are rufous. Head subglobose, as broad as long and about as thick as long, sparsely punctulate, with scattered white pilosity; occipital border rounded, sides in front of eyes very slightly concave. Mandibles thick and short; bifid at tip. Eyes large, orbicular, little convex, black, located at sides of head posterior to middle. Ocelli small, arranged in a triangle. Antennæ inserted at bases of flattened tubercles; contiguous at base; the scape clavate, about as long as head; apex of scape shortly excavated beneath, unarmed. Pedicel thickened toward apex, as long as the first funicular joint; first funicular joint a little longer than the second, joints 2-6 subequal; the last four subequal, forming a club. Prothorax above very short; in profile triangular; the pronotum thickly covered with white pubescence. Parapsidal furrows shallow, but distinct, extending the length of mesonotum. Mesonotum feebly, very sparsely punctate. Scutellum sparsely rugulose longitudinally, transversely



Fig. 3. *Hemilexis jessei* sp. nov. ♀

depressed posteriorly. Metanotum longitudinally carinate at middle, the surface rugulose. Petiole cylindrical, twice as long as broad, smooth and shining. Abdomen smooth and very shining; somewhat depressed. Legs rather short, the femora narrow basally, and clavate at apex. Wings extending past tip of abdomen, the tips rounded; pilose; slightly infuscated; veins yellowish.



Described from a series found in several colonies of *Formica subcyanea* Wheeler about Velasco and near Guerrero Mill. This is the second American species of *Hemilexis* as defined by Kieffer (*Gen. Ins. Fasc.* 124, p. 20, 1911), and differs widely from the other, *H. californica* Ashmead. Nothing is known regarding the habits of the other species, but *jessei* and the following variety are undoubtedly closely associated with the ants, probably as parasites on the immature phases. Only one or two individuals were found in each nest. They moved about rather slowly among the ants, which paid no attention to them.

This species is dedicated to my small collecting companion, Master Jesse Van Law.

*Hemilexis jessei* var. *minor* var. nov.

Three females which were taken in the same locality and with the same host ant as the preceding form seem to belong to a distinct variety. The size is much smaller (length 1.75 mm.) than in the typical form, and the coloration is lighter. The prothoracic pleuræ, scutellum and petiole being light fuscus. Otherwise the two forms are identical.

Family BETHYLIDÆ.

*Bruesiella* gen. nov.

Female. Apterous. Head transverse, broader than thorax; face evenly rounded, smooth except small fovæ at apex. Antennæ 12-jointed, the basal joint much enlarged; inserted at base of clypeus; narrowly separated at base. Mandibles stout, thickened at tips. Eyes well developed, oval, flat. Ocelli absent. Prothorax transverse, smooth, with a single large puncture near a feeble longitudinal impression at margin. Mesothorax transverse, the pleuræ extended into a thick, nearly perpendicular lamina. Metanotum from above subquadrate, broadest behind, sides feebly, irregularly margined; the posterior border with an acute, interrupted carina. Scutellum absent. Abdomen slender, somewhat depressed. Legs short, stout, the middle and posterior femora swollen, flattened; tibiæ spinose, the middle more strongly than the others.

This genus runs out at *Sclerochroa* in Kieffer's table in the *Genera Insectorum*. (Fascicle 76, p. 10, 1908). The peculiar structure of the mesothorax is quite different from that in any other of the described allied genera. In general appearance it bears considerable resemblance to some of the *Thynnidæ*.

*Type: Bruesiella formicaria* sp. nov.

*Bruesiella formicaria* sp. nov.

*Female:* (Fig. 4). Length 3.5 mm. Body shining. Head, thorax, anterior portion of first abdominal segment and legs brown, remainder of abdomen black.